

Mouse Monoclonal Antibody to P16 (Mouse and Human)

	Order I	nformation
Catalog#	20129	
Size/Concentration	100µl	50µl
Price(¥)	2180	1280

Description

The progression of cells through the cell cycle is regulated by a family of protein kinases known as cyclin-dependent kinases (Cdks). The sequential activation of individual members of this family and their consequent phosphorylation of critical substrates promotes orderly progression through the cell cycle. The cyclins function as differentially expressed positive regulators of Cdks. Negative regulators of the cycle include the p53-inducible 21 kDa WAF1/Cip1 protein designated p21, Kip1 p27 and p16. The complexes formed by Cdk4 and the D-type cyclins have been strongly implicated in the control of cell proliferation during the G1 phase. It has recently been shown that p16 binds to Cdk4 and inhibits the catalytic activity of the Cdk4/cyclin D complex. Moreover, the gene encoding p16 exhibits a high frequency of homozygous deletions and point mutations in established human tumor cell lines.

Specification		
Aliases : P16		
Entrez GeneID : 1029		
Swissprot : P42771		
clone : 2D9A12		
Host/Isotype : Mouse IgG2b		
Storage : Store at 4°C short term. Aliquot and store at -20°C long term. Avoid		
freeze/thaw cycles.		
Species Reactivity : Human,Rat		
Immunogen : Purified recombinant fragment of P16 expressed in E. Coli.		
Formulation : Ascitic fluid containing 0.03% sodium azide.		
Application		
IHC 1/200 - 1/1000		

IHC

ELISA

References

1/10000

1. Hunter, T. 1993. Cell 75: 839-841.


2. Sherr, C.J. 1993. Cell 73: 1059-1065.

3. El-Deiry, W.S., et al. 1993. Cell 75: 817-825.

Call 1-510-860-4615 +86-19375157864 Email Info@ProMab.com Web www.ProMab.com www.ProMab.cn

Protocal

WB - www.promab.com/protocol/wb.html IHC - www.promab.com/protocol/ihc.html ICC - www.promab.com/protocol/icc.html HCM - www.promab.com/protocol/hcm.html Antigen Sequence is available upon request.




Products and Services





Mouse Monoclonal Antibody to P16 (Mouse and Human)



Immunohistochemical analysis of paraffinembedded rat liver tissue (A), human brain tissue (B) and brain tumor (C), showing nuclear localization using P16 mouse mAb with DAB staining.

