

Mouse Monoclonal Antibody to NCOA3

	Order In	nformation
Catalog#	20274	
Size/Concentration	100µl	50µl
Price(¥)	2180	1280

Description

NCOA3: nuclear receptor coactivator 3. The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Two transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein.

Specification

Aliases : ACTR; AIB1

Entrez GeneID : 8202

Swissprot : Q9Y6Q9

clone : 2C11B12

ELISA

Host/Isotype : Mouse IgG1

Storage : Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Species Reactivity : Human

Immunogen : Purified recombinant fragment of NCOA3 (aa1-200) expressed in E. Coli.

Formulation : Ascitic fluid containing 0.03% sodium azide.

Application

1/10000

References

1. Mol Cell Biol. 2005 Sep;25(18):8273-84.


2. J Clin Oncol. 2006 Oct 1;24(28):4565-9.

Call 1-510-860-4615 +86-19375157864 Email Info@ProMab.com Web www.ProMab.com www.ProMab.cn

Protocal

WB - www.promab.com/protocol/wb.html IHC - www.promab.com/protocol/ihc.html ICC - www.promab.com/protocol/icc.html HCM - www.promab.com/protocol/hcm.html **Antigen Sequence** is available upon request.

Products and Services



For Research Only Application Key:WB - Western Blot | IHC - Immunohistochemistry | ICC - Immunocytochemistry | FCM -Flow Cytometry | ELISA - Enzyme-linked Immunosorbent Assay | IP - Immunoprecipitation

