

Mouse Monoclonal Antibody to NQO1

	Order Iı	nformation
Catalog#	30293	
Size/Concentration	100µl	50µl
Price(¥)	2180	1280

Description

This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. This FAD-binding protein forms homodimers and reduces quinones to hydroquinones. This protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Specification

Aliases : DTD; QR1; DHQU; DIA4; NMOR1; NMORI

Entrez GeneID : 1728

Swissprot : P15559

clone : 1A11

WB Predicted band size : 31kDa

Host/Isotype : Mouse IgG1

Storage : Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Species Reactivity : Human

Immunogen : Purified recombinant fragment of human NQO1 expressed in E. Coli. Formulation : Ascitic fluid containing 0.03% sodium azide.

Ap	plication	
	1/500 - 1/2000)

1/200 - 1/400

1/10000

FCM ELISA

WB

References

1. Mol Cancer Ther. 2009 Dec;8(12):3369-78.


2. J Biol Chem. 2009 Nov 27;284(48):33233-41.




Protocal

WB - www.promab.com/protocol/wb.html IHC - www.promab.com/protocol/ihc.html ICC - www.promab.com/protocol/icc.html HCM - www.promab.com/protocol/hcm.html **Antigen Sequence** is available upon request.


Products and Services




#30293




Mouse Monoclonal Antibody to NQO1



Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);



Western blot analysis using NQO1 mouse mAb against A549 (1), Hela (2), MCF-7 (3) and HepG2 (4) cell lysate.



Flow cytometric analysis of HepG2 cells using NQO1 mouse mAb (green) and negative control (red).

